



Europäisches Patentamt

European Patent Office

Office européen des brevets



⑪ Publication number : 0 596 684 A1

⑫

EUROPEAN PATENT APPLICATION

⑪ Application number : 93308719.9

⑫ Date of filing : 01.11.93

⑬ Int. Cl.⁵: C07C 37/045, C07C 67/29,
C07C 253/00, C07C 39/07,
C07C 39/27, C07C 39/28,
C07C 41/26, C07C 43/23,
C07B 41/02, C07C 205/22

⑭ Priority : 05.11.92 GB 9223195

⑮ Date of publication of application :
11.05.94 Bulletin 94/19⑯ Designated Contracting States :
AT BE CH DE DK ES FR GB GR IE IT LI LU MC
NL PT SE⑰ Applicant : OCTEL CHEMICALS LIMITED
Halebank
Widnes, Cheshire WA8 8NS (GB)

⑱ Inventor : Gilbert, Bruce Charles
58 Heworth Village
York, YO3 OAL (GB)
Inventor : Hanson, Peter
58 Moorgate, Acomb
York, YO2 4HR (GB)
Inventor : Jones, Jason Robert
168 Armadale Road
Ladybridge, Bolton, Lancs, BL3 4TP (GB)
Inventor : Timms, Allan William
5 Warrington Avenue, Whitby
Ellesmere Port, South Wirral, L65 6TR (GB)

⑲ Representative : Cardwell, Stuart Martin
Roystons Tower Building Water Street
Liverpool, Merseyside L3 1BA (GB)

⑳ Preparation of substituted aryl compounds.

㉑ A process for the preparation of aromatic compounds of the general formula I,



in which Ar represents an optionally substituted aryl group and X represents a group OH, Cl, Br or CN,
by reaction of an aryl diazoniumsalt of the formula II,



in which Ar is as defined above and Y represents a suitable counterion, with water, hydrogen chloride or
a salt thereof, hydrogen bromide or a salt thereof, or a cyanide in the presence of a copper(I)salt,
characterised in that the copper(I)salt is a copper(I)salt derived from an hydroxy carboxylic acid.

EP 0 596 684 A1

The present invention relates to a process for the preparation of aromatic compounds of the general formula I,



in which Ar represents an optionally substituted aryl group and X represents a group OH, Cl, Br or CN, by reaction of an aryl diazonium salt of the formula II,



in which Ar is as defined above and Y represents a suitable counterion, with water, hydrogen chloride or a salt thereof, hydrogen bromide or a salt thereof, or a cyanide in the presence of a copper(I) salt. The invention especially relates to the production of substituted phenols from the corresponding anilines by means of diazotisation and hydrolysis. In particular the preparation of 4-fluorophenol is described, a difficult reaction to carry out using conventional techniques.

The preparation of aryl chlorides, aryl bromides and aryl cyanides from the corresponding aryl diazonium salts is well known in the literature. The reaction is usually indicated as the Sandmeyer reaction. The reaction is often carried out by heating aryl diazonium salts with cuprous chloride, cuprous bromide or cuprous cyanide in the presence of hydrochloric acid, hydrobromic acid or an alkali metal salt of hydrogen cyanide respectively.

Also the preparation of phenols by hydrolysis of benzenediazonium ions is well known in the literature. It involves the preparation of a diazonium salt, e.g. the diazonium hydrogensulphate by treatment of the aniline hydrogensulphate with sodium nitrite in dilute aqueous sulphuric acid, followed by hydrolysis in more concentrated aqueous sulphuric acid. The temperature of the hydrolysis is maintained at the boiling point of the aqueous acid (by proper adjustment of the concentration of the sulphuric acid) and the phenol formed removed from the reaction medium by means of steam distillation in order to minimise the coupling of the formed phenol with more diazonium salt.

In the production of 4-fluorophenol from 4-fluoroaniline a major problem arises with corrosion of the glass lined vessels by hydrofluoric acid. The hydrofluoric acid is considered to arise by nucleophilic displacement of fluoride from the 4-fluorobenzenediazonium hydrogensulphate. This is not surprising since the diazonium group is one of the most electron withdrawing groups known thereby activating the 4-fluoro group to nucleophilic aromatic substitution. Furthermore, the resulting 4-hydroxybenzenediazonium hydrogensulphate may itself be hydrolysed to hydroquinone and thence by reaction with further diazonium ions to "diazo tars".

The liberation of hydrofluoric acid and production of "Diazo Tars" is of course exacerbated by the high temperatures employed in the reaction. A method which avoids these high temperatures and the liberation of hydrofluoric acid is therefore highly desirable.

It is well known that copper salts may be utilised in the Sandmeyer reaction of substituted anilines to the corresponding chlorides (using copper chloride), bromides (using copper bromide), nitriles (using copper cyanide) and phenols (using copper sulphate). The copper halide or pseudohalide is normally present in the copper (I) oxidation stage. Copper (II) sulphate may be used for the preparation of phenols, aryl chlorides, aryl bromides and aryl cyanides, together with small amounts of a reducing agent such as ascorbic acid to initiate a catalytic cycle. The mechanism of the first stage of the reaction is considered to be the reduction of copper (II) to copper (I) by the reducing agent. The resulting copper (I) subsequently reduces the diazonium ion to produce a phenyl radical and nitrogen. Water ligated to copper (II) is believed to be transferred to the phenyl radical producing the phenol and regenerating copper (I). Thus once the reaction has been initiated the cycle should become truly catalytic in copper.


Unfortunately, the ligand transfer of water from copper (II) to the radical is not efficient and large amounts of copper sulphate have to be utilized. Thus what is in principle a catalytic cycle is not in practice. This has major drawbacks in industrial production both in terms of expense of reagent and effluent treatment costs.

It has now been found that the use of a copper salt of an hydroxy carboxylic acid dramatically improves the preparation of compounds of the general formula I as defined hereinbefore. Organic acids such as citric or tartaric acid are suitable. The amount of copper needed may be significantly reduced. The salt of the copper may be prepared in situ.

The present invention therefore relates to a process for the preparation of aromatic compounds of the general formula I,



in which Ar represents an optionally substituted aryl group and X represents a group OH, Cl, Br or CN, by reaction of an aryl diazonium salt of the formula II,



in which Ar is as defined above and Y represents a suitable counterion, with water, hydrogen chloride or a salt thereof, hydrogen bromide or a salt thereof, or a cyanide in the presence of a copper(I) salt, characterised in that the copper(I) salt is a copper(I) salt derived from an hydroxy carboxylic acid.

Suitable hydroxy carboxylic acid to be used in the process of the present invention are all organic com-

pounds carrying at least one carboxylic acid group and one hydroxy group. The hydroxy carboxylic acid may contain one or more substituents, e.g. aryl groups, halogen atoms, nitro groups, alkoxy groups, phenoxy groups etc., provided that the substituents do not interfere with the reaction.

5 Preferably the hydroxy carboxylic acid is a straight or branched carboxylic acid having from 2 to 20 carbon atoms, containing up to four carboxylic acid groups and containing up to four hydroxy groups. More preferably the hydroxy carboxylic acid is a mono, di or tri carboxylic acid having up to 12 carbon atoms and containing up to three hydroxy groups. Still more preferably the hydroxy carboxylic acid is mono or di carboxylic acid having up to 8 carbon atoms, such as tartaric acid, citric acid, malic acid, citramalic acid, lactic acid or gluconic acid, citric acid being especially preferred.

10 Suitable salts of hydrogen chloride, hydrogen bromide and hydrogen cyanide are water soluble metal salts, especially alkali metal and earthalkali metal salts, preferably alkali metal salts such as sodium and potassium salts.

Suitable groups Ar in the compounds of the general formula I comprise hydrocarbocyclic compounds comprising up to five rings, usually up to four rings, more usually up to three rings. Other suitable aryl groups are 15 heterocyclic groups, especially 5- or 6- membered heterocyclic groups containing one to four of the same or different heteroatoms, especially one or two, such as nitrogen, oxygen and sulphur. Especially suitable are thiophene, pyridine, pyrrole, furan and pyrazole. All these groups may be substituted by one or more substituents. In principle, every known substituent in organic chemistry may be used provided that the groups do not interfere with the reaction. Preferably the optionally substituted aryl group is a phenyl or naphthyl group or a 5- or 20 6- membered heterocyclic group, optionally substituted by one or more halogen atoms or nitro, cyano, alkyl, haloalkyl, alkoxy, haloalkoxy, optionally substituted amino, formyl, alkoxycarbonyl, carboxyl, phenyl or halo- or dihalo-phenyl groups. More preferably the optionally substituted aryl group is a phenyl group, optionally substituted by one or more fluorine, chlorine or bromine atoms or nitro, cyano, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, optionally substituted amino, formyl, C₁₋₆ alkoxycarbonyl, carboxyl, phenyl or halo- or dihalo-phenyl groups.

25 The group Y in the general formula II is suitably a halide atom or a hydrogensulphate group, a phosphate group, a tetrafluoroborate group or a hexafluorophosphate group. Preferably Y represents a chloride or bromide atom or a hydrogensulphate group. In the case that an aryl chloride is to be prepared Y is preferably a chlorine atom, in the case that an aryl bromide is to be prepared Y is preferably a bromide, and in the case a phenol is to be made Y is preferably a hydrogensulphate.

30 A particular preferred process according to the present invention is the preparation of aromatic compounds of the general formula I in which X represent an OH group, by reaction of an aryl diazonium salt of the general formula II with water.

35 The copper(I)salts to be used in the process according to the present invention are suitably commercially available compounds as Cu Cl, Cu Br and Cu CN. Cuprous chloride will usually be used for the preparation of aromatic chlorides, cuprous bromide for the preparation of aromatic bromides, and cuprous cyanide for the preparation of aromatic cyanides. Instead of using copper(I)salts it is also possible to start with copper(II)salts, followed by in situ reduction to copper(I)salts. The invention therefore also relates to the processes as described hereinbefore, in which the copper(I)salt is prepared by reduction of a copper(II)salt in situ. Suitable reducing 40 agents are ascorbic acid, ferrous sulphate, hydrazine, bisulphite and titanium(III)chloride. Preferably ascorbic acid is used.

In a preferred embodiment of the invention a copper(II)salt is also present in the reaction mixture. The presence of the copper(II)salts results in improved yields.

45 In another preferred embodiment of the reaction the reaction is carried out in the presence of a water immiscible organic solvent in which the product of the reaction will dissolve. Suitable water immiscible solvents are alkanes as pentane, hexane, heptane, octane, cyclohexane and methylcyclohexane, halogenated alkanes as dichloromethane, chloroform, tetrachloromethane, trichloroethane and trichloroethene, aromatic solvents as benzene, toluene and xylene, halogenated aromatic solvents as chlorobenzene, ketones as methyl t-butyl ketone, and esters as methyl acetate, ethyl acetate and butyl acetate. A preferred water immiscible solvent is 50 ethyl acetate.

The process of the present invention may be carried out at a temperature between 0 °C and 100 °C. It is preferred to carry the reaction out at a temperature between 15 °C and 25 °C. In the case of the preparation of 4-fluorophenol no fluoride is liberated from the diazonium ion at that temperature.

55 It will be appreciated that the invention also relates to aromatic compounds of the general formula 1 as defined hereinbefore whenever prepared by a process as described above.

The invention will now be described in more detail by the following examples.

EXAMPLE 1

a. 4-Fluorobenzenediazonium hydrogensulphate

5 4-Fluoroaniline (11.2g; 0.1 mol) was added dropwise into a solution of sulphuric acid (25 g) in water (60 ml) over 10 min. A smooth paste was formed. The mixture was cooled to $<5^{\circ}$ and a solution of sodium nitrite (7.8 g; 0.11 mol) in water (25 ml) was added dropwise over 0.5 hours while maintaining the temperature at $<5^{\circ}$ with external cooling. A clear red solution was obtained. The solution was stirred for 0.25 hours to complete the reaction and urea (1.0 g) was added to destroy the excess nitrous acid. The resulting solution was used
10 directly in the next stage.

b. 4-Fluorophenol

Copper sulphate pentahydrate (6.25g; 0.025 mol) and trisodium citrate dihydrate (14.7 g; 0.05 mol) were
15 dissolved in water (60 ml). Ethyl acetate (150 ml) was added and the two phase mixture vigorously stirred. A few ml of the diazonium solution prepared above was added to the stirred mixture. The reaction was initiated by the addition of a solution of ascorbic acid (1.0 g) dissolved in water (20 ml). The remaining solution had been added, the mixture was allowed to settle and the organic phase separated. The aqueous phase was re-extracted with more ethyl acetate and the combined organic phases washed with a little water. The solvent was
20 removed by evaporation under reduced pressure and the residue purified by means of distillation. The product (9.0 g; 80%) was obtained by b.p. $81^{\circ}/13$ mm. Hg. Fluoride ion in the aqueous liquors could not be detected.

EXAMPLES 2-12

25 Similarly prepared by the methods cited in Example 1 the following compounds are prepared

Example	Substituent	b.p. (m.p.)
2	4-Chloro	(41-43 $^{\circ}$)
3	4-Methoxy	(55-57 $^{\circ}$)
4	4-Methyl	201-202 $^{\circ}$
5	Hydrogen	181-183 $^{\circ}$
6	4-Nitro	(112-114 $^{\circ}$)
7	4-Cyano	(111-113 $^{\circ}$)
8	4-Ethoxycarbonyl	(116-118 $^{\circ}$)
9	2-Fluoro	171-173 $^{\circ}$
10	2-Chloro	175-176 $^{\circ}$
11	2-Methyl	202-204 $^{\circ}$
12	2-Methoxy	204-206 $^{\circ}$

EXAMPLE 13 (for comparative purposes only)

50 4-Chlorobenzonitrile

Sodium cyanide (2.94g; 0.06 mol) was added to water and stirred. Cuprous cyanide (2.7g; 0.03 mol) was added to the mixture stirred until it became homogeneous. 4-chlorobenzenediazonium tetrafluoroborate (5.66g; 0.025 mol) in water (10 ml) was added to one portion. The mixture was allowed to react at room temperature. The mixture was extracted by ethyl acetate and the extract analysed by means of gas chromatography which indicated that 4-chlorobenzonitrile had been formed in 75% yield.
55

EXAMPLE 14

4-Chlorobenzonitrile

- 5 The method of example 13 was carried out except that trisodium citrate (0.06 mol) was added after the copper cyanide but before the diazonium salt. The reaction was conducted for 1 hour when a yield of 90.4% was obtained by analysis of the ethyl acetate extract.

EXAMPLE 15

10

4-Chlorobenzonitrile

- 15 Copper (II) nitrate (0.03 mol) and trisodium citrate (0.06 mol) were dissolved in water and the pH adjusted to 9 - 10 with sodium hydroxide solution. Sodium cyanide (0.06 mol) was added and the mixture stirred. 4-Chlorobenzenediazonium tetrafluoroborate (0.025 mol) was added in one portion. After 10 mins the reaction was adjudged to be complete and analysis by gas chromatography indicated a yield of >90%.

Claims

20

1. A process for the preparation of aromatic compounds of the general formula I,



in which Ar represents an optionally substituted aryl group and X represents a group OH, C 1, Br or CN, by reaction of an aryl diazonium salt of the formula II,

25



in which Ar is as defined above and Y represents a suitable counterion, with water, hydrogen chloride or a salt thereof, hydrogen bromide or a salt thereof, or a cyanide in the presence of a copper(I)salt, characterised in that the copper(I)salt is a copper(I)salt derived from hydroxy carboxylic acid.

- 30 2. A process according to claim 1, in which the hydroxy carboxylic acid is a straight or branched carboxylic acid having from 1 to 20 carbon atoms, containing up to four carboxylic acid groups and containing up to four hydroxy groups.
3. A process according to claim 2, in which the hydroxy carboxylic acid is a mono, di or tri carboxylic acid having up to 12 carbon atoms and containing up to three hydroxy groups.
- 35 4. A process according to claim 1, in which the hydroxy carboxylic acid is tartaric acid, citric acid, malic acid, citriamalic acid, lactic acid or gluconic acid.
- 40 5. A process according to claim 1, in which the optionally substituted aryl group is a phenyl, naphthyl or 5- or 6- membered heterocyclic group optionally substituted by one or more halogen atoms or nitro, cyano, alkyl, haloalkyl, alkoxy, haloalkoxy, optionally substituted amino, formyl, alkoxy carbonyl, carboxyl, phenyl or halo- or dihalo-phenyl groups.
- 45 6. A process according to claim 5, in which the optionally substituted aryl group is a phenyl group, optionally substituted by one or more fluorine, chlorine or bromine atoms or nitro, cyano, C₁₋₆ alkyl, C₁₋₆ Haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, optionally substituted amino, formyl, C₁₋₆ alkoxy carbonyl, carboxyl, phenyl or halo- or dihalo-phenyl groups.
- 50 7. A process as claimed in any of claims 1 to 6, in which Y represents a chloride or bromide atom or a hydrogensulphate group.
8. A process as claimed in claim 7, in which Y represents a hydrogensulphate group.
- 55 9. A process as claimed in any of claims 1 to 8, in which an aromatic compound of the general formula I in which X represents an OH group, is prepared by reaction of a aryl diazonium salt of the general formula II with water.
10. A process as claimed in any of claims 1 to 9, in which the copper(I)salt is prepared by reduction of a cop-

per(II)salt in situ.

- 5
11. A process as claimed in claim 10, in which the reducing agent is ascorbic acid, ferrous sulphate, hydrazine, bisulphate or titanium(III)chloride.
12. A process as claimed in any of claims 1 to 11, in which also a copper(II)salt is present in the reaction mixture.
13. A process as claimed in any of claims 1 to 12, in which the reaction is carried out in the presence of a water immiscible organic solvent.
- 10
14. A process as claimed in claim 13, in which the water immiscible solvent is ethyl acetate.
15. A process as claimed in any of claims 1 to 12, in which the reaction is carried out at a temperature between 0°C and 100°C.
- 15
16. A process as claimed in claim 15, in which the temperature is between 15°C and 25°C.
17. Aromatic compounds of the general formula 1 as defined in any of the preceding claims whenever prepared by a process according to any of claims 1 to 16.
- 20
- 25
- 30
- 35
- 40
- 45
- 50
- 55

European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 93 30 8719

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.-CL.5)
A	FR-A-2 234 260 (OLIN CORPORATION) * the whole document *	1	C07C37/045 C07C67/29 C07C253/00 C07C39/07 C07C39/27 C07C39/28 C07C41/26 C07C43/23 C07B41/02 C07C205/22
A	CHEMICAL ABSTRACTS, vol. 107, no. 3, 20 July 1987, Columbus, Ohio, US; abstract no. 23078y, page 585 ;column 2 ; * abstract * & JP-A-62 029 544 (NIPPON KAYAKU CO.)	1	
A	DATABASE WPI Derwent Publications Ltd., London, GB; AN 75-68040W & JP-A-50 076 038 (DAINI SEIKOSHA) * abstract *	1	
			TECHNICAL FIELDS SEARCHED (Int.-CL.5)
			C07C C07B
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 28 January 1994	Examiner Heywood, C
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1800 03.12 (P04C01)